

## 2018-2019 Genetic Written PT

- 1.) True/False: The color intensity in ELISA is proportional to the amount of conjugate bound.
- 2.) True/ False:: The acceptable MINIMUM number of seeds used for a germination test is 500.
- 3.) True/ False: In a herbicide bioassay test, if the positive trait check sample is not growing or the negative trait check sample is growing normally, this is a sign that the test is proceeding as it should.
- 4.) True/False: During an ELISA test, the purpose of a wash buffer is to remove unbound materials and keep bound components stable.
- 5.) True/ False: Though antibodies are usually very specific, the antibody for the Soybean Mosaic Virus will bind to the Bt-Cry1Ab antigen.
- 6.) True/False: Pipette tips used in ELISA can be reused if you rinse them out first.
- 7.) True/False: ELISA substrates are sensitive to light and exposure to light must be minimized.
- 8.) True/False In herbicide bioassays, the check samples must be exposed to the same conditions as each replicate from the test sample.
- 9.) Herbicides can be formulated as all of the following except:
  - a. Dry Flowable
  - b. Dry Granules
  - c. Compressed Gas
  - d. Liquid
- 10.) Which herbicide bioassay method involves adding the herbicide to water that moistens the medium and placing the seeds on the medium, allowing them to sit for 7 days?
  - a. Presoak Method
  - b. Substrate Imbibition
  - c. Seedling Spray
- 11.) In herbicide bioassays, test sample growth could be irregular for all the following reasons except:
  - a. Seed quality is low
  - b. Herbicide concentration is too high
  - c. Wrong check sample was used
  - d. Test sample is not tolerant to the herbicide
- 12.) Which herbicide bioassay method involves an herbicide pretreatment before the seeds are placed on the germination medium?
  - a. Presoak Method
  - b. Substrate Imbibition
  - c. Seedling Spray

## 2018-2019 Genetic Written PT

- 13.) In which type of ELISA is the antigen coated directly to the plate?
- Double Antibody Sandwich (DAS) ELISA
  - Compound Direct ELISA
  - Compound Indirect ELISA
- 14.) The values given for each well by a microplate spectrophotometer during quantitative ELISA represent:
- Optical density
  - Concentration in ug/mL
  - Volume
- 15.) The first seed soak method for herbicide bioassay was used to select for herbicide tolerant plants on which crop?
- Sulfonylurea tolerant soybeans (STS)
  - Imidazolinone tolerant corn (IMI)
  - Buctryl resistant cotton
  - Roundup Ready cotton
- 16.) Which part of "ELISA" refers to the antibody-antigen interaction?
- Enzyme
  - Linked
  - Immuno
  - Sorbent
  - Assay
- 17.) What is the minimum number of seeds that need to be planted to report results at a 99% confidence level in herbicide bioassay?
- 300
  - 400
  - 500
  - 600
- 18.) In ELISA, what explanations should be considered for no color present in the wells? (select all that apply)
- wrong antibodies used
  - wrong enzyme conjugate used
  - substrate not added
  - wrong substrate used
  - wrong buffers used
  - enzyme conjugated not added

## 2018-2019 Genetic Written PT


19.) Select the advantages of substrate imbibition. (select all that apply)

- A. automation of method,
- B. ease of including check sample with each replicate,
- C. less steps to plant
- D. results available in 24 hours

20.) What are the advantages of the seedling spray method? (select all that apply)

- a. relates well to field conditions
- b. turnaround time
- c. seed quality is of less concern since only normal seedlings are sprayed
- d. Cheapest method with little resources used.

21.) Labels the antigen-positive lateral flow strip with the following areas: wick area, sample verification area, antibody specific area, sample uptake area.



22.) How many grams of Lightning would you need to make one liter of 100 ppm stock solution from a 70% a.i. Lightning solution? Show your work.

23.) 4 normal non-trait seedlings, 2 abnormal seedlings, and 394 normal trait seedlings were obtained in a 400 seed test. What percentage of seedlings contained the desired herbicide trait?

## 2018-2019 Genetic Written PT

24.) Which of the two most common ELISA enzyme conjugates is present in human and animal bodily fluids?

25.) Name three symptoms non-trait soybeans could express when exposed to an herbicide.

26.) What is the difference between how polyclonal antibodies and monoclonal antibodies are produced?

27.) Place the following components in order of reaction for use in Double Antibody Sandwich (DAS) ELISA. Enzyme conjugate, antibody, substrate, sample or antigen

28.) Name five crops that use herbicide traits.

29.) Name three symptoms non-trait corn could express when exposed to an herbicide.

30.) What was the initial purpose of ELISA upon creation of the method?

31.) You run an ELISA plate and the plate reader gives you the following optical density results for the first column – 0.984, 0.769, 0.039, 0.844, 0.046, 0.945, 0.052, & 0.799. How many would you consider as having a negative result?